

A new genus, *Smilirhexia*, of Smiliini (Hemiptera, Membracidae) from Costa Rica

Stuart H. McKamey

Systematic Entomology Laboratory, Plant Sciences Institute, Agricultural Research Service, USDA, c/o National Museum of Natural History, P.O. Box 37012, Smithsonian Institution, Washington, D.C.

urn:lsid:zoobank.org:author:E6BD01B8-0684-464A-8A86-812701DA05C7

Corresponding author: *Stuart H. McKamey* (stuart.mckamey@ars.usda.gov)

Guest editor: *Lewis Deitz* | Received 17 September 2008 | Accepted 6 October 2008 | Published 31 October 2008

urn:lsid:zoobank.org:pub:45774FDF-CF37-4A92-A63F-8A0BFC15B5A9

Citation: McKamey SH (2008) A new genus, *Smilirhexia*, of Smiliini (Hemiptera, Membracidae) from Costa Rica. ZooKeys 3: 51-55. doi: 10.3897/zookeys.3.29

Abstract

Smilirhexia naranja, **gen. n.** and **sp. n.**, is described from Central America, the southern limit of the tribe Smiliini, and represents a strong divergence from the morphology of the oak-feeding genera prevalent in North America.

Keywords

Membracoidea, Homoptera, Auchenorrhyncha, new genus, new species

Introduction

The tribe Smiliini is the nominotypical tribe of Smiliinae, one of the largest subfamilies of treehoppers (McKamey 1998). The subfamily is native to the New World, being most speciose in the tropics. In contrast to this subfamilial pattern, the species of Smiliini are primarily North American, with the great majority of species oligophagous on oaks (*Quercus* spp., Fagaceae) and the southernmost species recorded from Panama (R. Cocroft, pers. comm.). Excluding the odd smiliine genus *Antianthe* Fowler, the tribe's southern limit is even more restricted than that of oaks, which extend southward to Colombia.

The mostly recently described species is *Telamona archiboldi* Froeschner (1968), from Florida. There have been only three recent taxonomic works on Smiliini (McKamey and Deitz 1991, Wood and Pesek 1992, Wallace and Deitz 2003). The other

recent papers consist of local or regional checklists (Kopp and Yonke 1973, 1974; Johnson and Freytag 1997; Wallace et al. 2003; Flynn et al. 2003; Wallace and Troyano 2006). Quisenberry et al. (1978) described the immatures of most genera of Smiliini.

Morphological terminology follows Deitz (1975).

***Smilirhexia* McKamey, gen. n.**

urn:lsid:zoobank.org:act:AD148098-EBFC-46EA-BF5D-B74F697FD853

Type species: *Smilirhexia naranja*, sp. n.

Diagnosis. The new genus differs from all other genera of the tribe in having a smooth and shining pronotum, which is broad, almost completely convex, and brightly colored.

Description of male. Head (Fig. 3) glabrous, subtriangular in anterior view, ocelli equidistant from each other and from eyes. Pronotum (Figs 1-3) entirely concealing scutellum, convex, dorsal carina elevated only in posterior half, and weakly, otherwise delineated by absence of punctures; surface shining, punctures shallow and lacking setae; humeral angles not projected laterally beyond wing bases; posterior pronotal process ending at apex of forewing cell M_{3+4} (last apical cell). Forewing (Fig. 1) free, in repose not concealed wholly or in part by pronotum; without r-m crossveins, with veins R and M adjacent basally, weakly divergent soon after, R_{4+5} and M_{1+2} confluent for a short distance before apex, and strongly divergent at wing apex; R_{2+3} present as distinct branch of R; R_4 and R_5 separately joined to vein M, creating a small cell R_4 near center of wing. Hind wing with short r-m crossvein (holotype right wing) or veins R and M fused at single point (holotype left wing). Femora, tibiae, and tarsomeres lacking cucullate setae except metathoracic femur with two apically, metathoracic tibia with three rows (row 2 double), and metathoracic tarsomere I with 4 cucullate setae at apex. Metathoracic coxa and trochanter without apposed processes. Abdomen lacking fenestrae or mid-dorsal protuberances.

Female. Unknown.

Distribution. Costa Rica, Puntarenas Prov.

Material Examined. Holotype male of *Smilirhexia naranja*, sp. n.

Etymology. The new genus name is feminine and combines the name of its apparent tribe and the name of an unrelated genus, *Rhexia* Stål, which it resembles superficially [e.g., *Rhexia viridicollis* (Fowler), illustrated by Deitz (1983)].

Discussion. The distally confluent radial and medial veins, which diverge before the forewing apex, and the lack of processes on the metathoracic coxa and trochanter place the new genus firmly in the subfamily Smiliinae. In Deitz's key to the higher classification, the genus keys to the tribe Smiliini due to the presence of 2 m-cu crossveins. The only discrepancy is that in the new genus, the forewing is completely exposed rather than partially concealed. Most genera of the tribe are morphologically similar and authentic records of only a few species exist for Central America, the tribe's southern limit. The notable exception is the genus *Antianthe* Fowler, which is laterally compressed, strongly crested, and with pronounced humeral angles, and which occurs from the southern


USA to South America. *Smilirhexia* appears to represent another distinct morphological form for *Smiliini* that may help to elucidate the evolutionary origin of this tribe.

Species of the subfamily Darninae, which appears to be a close relative of *Smiliinae*, also have the forewing with two r-m crossveins, as in *Smiliini*. Interestingly, *Cymbomorpha* Stål (Cymbomorphini, Darninae) occasionally have the characteristic forewing venation of *Smiliinae*, with veins R and M confluent for a short distance before the apex, and double setation in metathoracic tibial row II. Both *Cymbomorpha* and *Eumela* Stål, the other cymbomorphine genus, and *Smilidarnis* Andrade, 1989 (a Neotropical, unplaced genus with characters of both Darninae – especially Cymbomorphini – and *Smiliinae*), exhibit the shining pronotal surface and completely exposed forewings of the new genus *Smilirhexia*.

***Smilirhexia naranja* McKamey, sp. n.**

urn:lsid:zoobank.org:act:F473B88C-4AF2-45CF-9F7D-964FBCDF7B3F

Description of male. Length including forewings in repose 8.0 mm, width between humeral angles 4.6 mm. Color. Head yellow-orange. Pronotum yellow-orange dorsally and posteriorly, yellow laterally except with large lateral indentation at mid-length black. Forewing fuscous at coriaceous base and transversally at distal crossveins, cells tinted yellow basally, hyaline distally. Male terminalia (Figs 4-6): lateral plate distinct,



Figs 1-6. *Smilirhexia naranja*, sp. n. 1-3, Lateral, dorsal, and anterior views of head, pronotum, wings, and legs (forewing anterior apices presumably rolled inward as a preservative side-effect). 4, Lateral view, cleared terminalia before dissection. 5-6. Aedeagus in lateral and anterior views, respectively. *aed*, aedeagus; *lp*, unarmed lateral plate.

long, lacking protuberances. Aedeagus narrow in anterior view, in lateral view U-shaped and widest at mid-length, anterior margin with many denticulae of increasing size ventrally; styles simple, distally recurved (Fig. 4), apex acute.

Female. Unknown.

Distribution. COSTA RICA: Puntarenas Prov.: Coto Brus, nr. San Vito, Estación Biológica Las Alturas, 8° 57'N, 82°50'W.

Material Examined. Holotype male, pinned, dissected, with labels "Est. Biol. Las Alturas, / 1500m., Coto Brus, Prov. / Puntarenas. Costa Rica, / F. Araya, 23 mar a 2 may / 1992, L-S 322500,591300", "COSTA RICA INBIO / CR1000 / 777014", and "HOLOTYPE / *Smilirhexia* / *naranja* McKamey". Deposited at the Instituto Nacional de Biodiversidad (INBio), Santo Domingo de Heredia, Costa Rica.

Etymology. The species epithet is a Spanish adjective for orange, the predominant pronotal color of the species.

Notes. In Costa Rica, most oaks occur in submontane and montane forests (1500-3200m) (Kappelle et al. 1992). In the Cordillera de Talamanca, which contains the type locality, there are at least eight oak species, with at least three reported from Coto Brus itself: *Quercus seemannii* Liebm., *Q. oocarpa* Liebm., and *Q. rapurahuensis* Pittier ex Trel. (Giddy 2008), and another (*Q. insignis* M. Martens & Gal.) at 1300m on the trail to Las Alturas (Instituto Nacional de Biodiversidad 2008). In short, it is easily possible that this new treehopper genus, like others in the tribe, feeds on *Quercus*.

Acknowledgments

I thank Carolina Godoy, formerly of the Instituto Nacional de Biodiversidad, for access to the collection and loan of the type specimen. L. L. Deitz, C.-P. Lin, N. Vandenberg, and M. Pogue provided helpful comments on an earlier draft of the manuscript.

References

- Andrade GS (1989) *Smilidarnis*, a new genus with two new species of Membracidae from South America (Homoptera). *Revista Brasileira de Zoologia* 6(4): 695-699.
- Deitz LL (1975) Classification of the higher categories of the New World treehoppers (Homoptera: Membracidae). North Carolina Agricultural Experiment Station Technical Bulletin 225: 1-177.
- Deitz LL (1983) Name changes in the Membracidae. *Proceedings of the Entomological Society of Washington* 85(4): 856-857.
- Flynn DJ, Balme GR, Deitz LL, Rothschild MJ (2003) *Archasia pallida* (Fairmaire) (Hemiptera: Membracidae) in North Carolina. *Proceedings of the Entomological Society of Washington* 105(4): 1060-1061.
- Froeschner RC (1968) *Telamona archboldi*, a new treehopper from Florida (Homoptera: Membracidae). *Proceedings of the Entomological Society of Washington* 70(2): 154-155.

- Giddy I (2008) Cloudbridge: Bridging a Costa Rican cloud forest. <http://cloudbridge.org> [accessed August 29, 2008]
- Instituto Nacional de Biodiversidad (2008) Lista de especímenes de *Quercus insignis*. <http://www.inbio.ac.cr/bims/k03/p13/c045/o0243/f01545/g007809/s023537.htm> [accessed August 29, 2008]
- Johnson MP, Freytag PH (1997) Treehoppers (Homoptera: Membracidae) on pin oak in Kentucky. *Journal of the Kansas Entomological Society* 70(1): 21-30.
- Kappelle M, Cleef AM, Chaverri A. (1992) Phytogeography of Talamanca montane *Quercus* forests, Costa Rica. *Journal of Phytogeography* 19: 299-315.
- Kopp DD, Yonke TR (1973) The treehoppers of Missouri: Part 3. Subfamily Smiliinae; tribe Smiliini. *Journal of the Kansas Entomological Society* 46(3): 375-421.
- Kopp DD, Yonke TR (1974) The treehoppers of Missouri: Part 4. Subfamily Smiliinae; tribe Telamonini (Homoptera: Membracidae). *Journal of the Kansas Entomological Society* 47(1): 80-130.
- McKamey SH (1998) Taxonomic catalogue of the Membracoidea (exclusive of leafhoppers). Second supplement to fascicle 1 – Membracidae of the general catalogue of the Hemiptera. *Memoirs of the American Entomological Institute* 60: 1-377.
- McKamey SH, Deitz LL (1991) Nomenclatural changes in the treehopper tribes Hoplophorini, Smiliini, and Talipedini (Homoptera: Membracidae). *Proceedings of the Entomological Society of Washington* 93(1): 193-196.
- Quisenberry SS, Yonke TR, Kopp DD (1978) Key to the genera of certain immature treehoppers of Missouri with notes on their host plants (Homoptera: Membracidae). *Journal of the Kansas Entomological Society* 51(1): 109-122.
- Wallace MS, Deitz LL (2003) *Aconophora lineosa* Walker 1858, new junior synonym of *Thelia bimaculata* (Fabricius 1794) (Hemiptera: Membracidae). *Proceedings of the Entomological Society of Washington* 105(4): 1051-1052.
- Wallace MS, Deitz LL, Rothschild MJ (2003) Checklist of the treehoppers (Hemiptera: Membracidae) of Great Smoky Mountains National Park. *Proceedings of the Entomological Society of Washington* 105(3): 578-591.
- Wallace MS, Troyano NM (2006) The oak-feeding smiliine treehoppers (Hemiptera: Membracidae) of Delaware Water Gap National Recreation Area. *Journal of the Pennsylvania Academy of Science* 80(1): 28-34.
- Wood TK, Pesek JD (1992) Pronotal shape: a source of confusion or panacea in systematic studies of treehoppers (Homoptera: Membracidae). In: Sorensen JT, Foottit R (Eds) *Ordination in the study of morphology, evolution and systematics of insects: applications and quantitative genetic rationals*. Elsevier, Amsterdam, London etc., 349-384.